

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</small>					
1. REPORT DATE (DD-MM-YYYY) 2006		2. REPORT TYPE Open Literature		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Growth factor dependent cholinergic function and survival in primary mouse spinal cord cultures				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Sheridan, R.E., Adler, M.				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: MCMR-CDT-N 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400				8. PERFORMING ORGANIZATION REPORT NUMBER USAMRICD-P05-004	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: MCMR-CDZ-P 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Published in Life Sciences, 79, 591-595, 2006					
14. ABSTRACT See reprint.					
15. SUBJECT TERMS Mouse, spinal cord, cell culture, growth factor, CNTF, BDNF, bFGF, choline, acetylcholine, glycine.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UNLIMITED	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON Michael Adler
a. REPORT UNCLASSIFIED	b. ABSTRACT UNCLASSIFIED	c. THIS PAGE UNCLASSIFIED			19b. TELEPHONE NUMBER (include area code) 410-436-1913



Growth factor dependent cholinergic function and survival in primary mouse spinal cord cultures[☆]

Robert E. Sheridan, Michael Adler^{*}

Neurobehavioral Toxicology Branch, Analytical Toxicology Division, U.S. Army Medical Research Institute of Chemical Defense, 3100 Ricketts Point Road, Aberdeen Proving Ground, MD 21010-5400, United States

Received 14 June 2005; accepted 1 February 2006

This article is dedicated to the memory of Dr. Robert E. Sheridan

Abstract

In primary embryonic spinal cord cultures, synaptic transmission can be conveniently studied by monitoring radiolabeled neurotransmitter release or by recording of electrophysiological responses. However, while the mature spinal cord contains an appreciable number of cholinergic motoneurons, cultures of embryonic spinal cord have a paucity of these neurons and release little or no acetylcholine upon stimulation. To determine whether the proportion of cholinergic neurons in primary mouse spinal cord cultures can be augmented, the effects of several classes of growth factors were examined on depolarization- and Ca^{2+} -evoked release of choline/acetylcholine (Ch/ACh). In the absence of growth factors, little or no evoked release of radiolabeled Ch/ACh could be demonstrated.

Media supplemented with brain-derived neurotrophic factor (BDNF), ciliary neurotrophic factor (CNTF) or basic fibroblast growth factor (bFGF) were examined for their ability to preserve the population of neurons in culture. CNTF was found to increase the number of surviving neurons and to enhance the release of radiolabeled Ch/ACh; the other factors were without effect. The action of CNTF was transient, and the neuronal population decreased to levels observed in cultures lacking growth factor after 20 days in vitro. The correlation between enhanced neuron survival and increased Ch/ACh release suggests that CNTF protected cholinergic neurons, albeit transiently, from cell death.

Published by Elsevier Inc.

Keywords: Mouse; Spinal cord; Cell culture; Growth factor; CNTF; BDNF; bFGF; Choline; Acetylcholine; Glycine

Introduction

Primary cultures of dissociated embryonic mammalian spinal cord represent a convenient system to study mechanisms of transmitter release. Radiolabeled neurotransmitters or precursors

are internalized by neurons in culture and released quantitatively in response to depolarization and Ca^{2+} ; this release is inhibited by tetanus toxin and botulinum neurotoxins in a concentration- and time-dependent manner (Bergey et al., 1987; Williamson et al., 1992; Keller et al., 2004; Hall et al., 2004; Sheridan et al., 2005). The sensitivity of spinal cord cultures to botulinum neurotoxin (BoNT) appears to be greater than that of other cell systems described, and spinal cord motoneuron terminals represent an important biological target for BoNT (Daniels-Holgate and Dolly, 1996; Simpson, 2004). Using these cultures, it has been possible to demonstrate a relationship between toxin-mediated cleavage of SNARE proteins and inhibition of transmitter release (Williamson et al., 1996; Keller et al., 1999, 2004), even though the cultures contain a large proportion of non-neuronal cells which lack regulated exocytosis (Brenneman et al., 1987). The majority of transmitter release studies have focused on depolarization-

[☆] Opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the Army or the Department of Defense. All research was conducted in compliance with the Animal Welfare Act and other Federal statutes and regulations relating to animals and adheres to principles stated in the Guide for the Care and Use of Laboratory Animals, National Research Council, 1996. The facilities where this research was conducted are fully accredited by the Association for Assessment and Accreditation of Laboratory Animal Care (AAALAC) International.

^{*} Corresponding author. U.S. Army Medical Research Institute of Chemical Defense, ATTN: MCMR-CDT-N, 3100 Ricketts Point Road, Aberdeen Proving Ground, MD 21010-5400, United States. Fax: +1 410 436 8377.

E-mail address: Michael.adler@amedd.army.mil (M. Adler).

dependent release of glutamate or glycine, neither of which is the normal target of BoNT in vivo (Simpson, 2004). Unless spinal cord neurons are cocultured with muscle or cervical ganglion cells, little or no cholinergic activity has been observed (Bennett et al., 1980; Kobayashi et al., 1987; Wang et al., 1990; Braun et al., 1997).

Spinal cord preparations usually include dorsal root ganglion (DRG) cells, which are important in stabilizing synapses between neurons and skeletal muscle (Kobayashi et al., 1987); however, DRG cells alone are not capable of maintaining the population of cholinergic neurons (Wang et al., 1990). During the first weeks after plating, large numbers of neurons undergo cell death, the majority of which appear to be cholinergic motoneurons (Ransom et al., 1977). It has been reported that brain derived neurotrophic factor (BDNF) can enhance the short term (2–5 days) survival of spinal cord motoneurons in vitro (Wong et al., 1993; Henderson et al., 1993; Hughes et al., 1993). Similarly, basic fibroblast growth factor (bFGF) and ciliary neurotrophic factor (CNTF) have been reported to support the population of motoneurons in culture (Hughes et al., 1993; Wong et al., 1993), although the results have been inconsistent.

The current study is an attempt to determine the ability of neurotrophic factors to promote the survival of cholinergic neurons in primary mouse embryonic spinal cord cultures. CNTF was found to increase the number of surviving neurons and to enhance the release of radiolabeled choline/acetylcholine (Ch/ACh), while the other factors were without effect. The action of CNTF was transient, being significant only during week 2 in culture. These results suggest that CNTF provides a limited window for studies on cholinergic neurons in primary mouse spinal cord cultures.

Materials and methods

Spinal cord cultures

Spinal cord with DRG were removed from embryonic Hsd:NIHS Swiss mice at gestation day E13–E14, dissociated in 0.125% trypsin and plated at a density of 4×10^5 cells/cm² in 12-well collagen-coated tissue culture plates. The cells were cultured at 37°C in 90% air/10% CO₂ in Eagle's Minimum Essential Media (MEM) with 5% horse serum and N3 supplement (Romijn et al., 1981; Wang et al., 1990) or in neurobasal media with B27 supplement and 2% horse serum (Brewer, 1995). Cell cultures were treated with 54 μM 5-fluoro-2'-deoxyuridine and 140 μM uridine 5–9 days after plating to inhibit glial cell division. Thereafter, cells were fed twice per week. Growth factors were added to the culture media 2 days after plating and reapplied at each media change.

Determination of neuronal cell density

To identify the neuronal population in culture, the media was aspirated and cells were washed twice with phosphate-buffered saline, fixed in 4% paraformaldehyde for 30 min at room temperature and stained for neuron specific enolase (NSE) using the method of Brenneman et al. (1987). NSE positive cells were counted under a phase contrast microscope at 160×. Cells from

20 fields were averaged, and 4 wells were counted for each condition.

Neurotransmitter release assay

Neurotransmitter release was detected by the method of Williamson et al. (1992) and is described in detail by Sheridan et al. (2005). The growth medium was removed, and the cells were incubated at 37°C for 20 min in a HEPES-buffered salt solution (HBSS) containing ³[H]-glycine, ³[H]-glutamate, ³[H]-glutamine or ¹⁴[C]-Ch, as appropriate. The composition of the HBSS was 135 mM NaCl, 3 mM KCl, 1 mM MgCl₂, 2 mM CaCl₂, 10 mM glucose, 10 mM HEPES/NaOH (pH 7.3) and 0.1% w/v bovine serum albumin. The cells were then washed briefly with HBSS and sequentially incubated for 5 min each in (1) Ca²⁺-free HBSS containing 0.5 mM EGTA, (2) HBSS containing 56 mM K⁺+2 mM Ca²⁺ and (3) Ca²⁺-free HBSS containing 0.5 mM EGTA. Each incubation solution was collected and the radioactivity was determined by scintillation counting. Following these incubations, the cells were lysed in 0.2% sodium dodecyl sulfate and the remaining radiolabel in the cells was determined. The labeled neurotransmitter selectively released with high (56 mM) K⁺ in the presence of 2 mM Ca²⁺ was expressed as a fraction of the total radioactivity in the cells. Cells incubated with ¹⁴[C]-Ch release ACh as well as unmodified Ch (McGee et al., 1978). Since we did not separate these products, the materials released after incubation with ¹⁴[C]-Ch are designated collectively as Ch/ACh in the present study.

Data analysis

Unless stated otherwise, all data are expressed as the mean ± SEM. Statistical analysis was performed using a one-way analysis of variance followed by Bonferroni's posttests (Graphpad InStat ver 3.0, San Diego, CA, USA). $P \leq 0.01$ was considered to be statistically significant.

Materials

MEM, Neurobasal media, B27, horse sera, 5-fluoro-2'-deoxyuridine and uridine were obtained from Invitrogen Corp. (Carlsbad, CA, USA). BDNF (human recombinant), CNTF (rat recombinant), and routine reagents were purchased from Sigma-Aldrich (St. Louis MO, USA). Bovine recombinant bFGF was obtained from Roche Applied Science (Indianapolis, IN, USA). Growth factors were applied at concentrations twice their EC₅₀, as provided by the supplier. Radiolabeled neurotransmitters and precursors were obtained from Perkin Elmer Life and Analytical Sciences (Shelton, CT, USA).

Results

Cell death in spinal cord cultures

Following the initial plating of mouse embryonic spinal cord cells, there was a substantial loss of neurons before the




Fig. 1. Effect of growth factors on neuronal survival. Spinal cord neurons were counted in 20 random 160× microscopic fields at the indicated times after plating under control conditions (●) or following treatment with bFGF (■, 2 ng/ml), BDNF (▲, 1 ng/ml) or CNTF (▼, 10 ng/ml). Growth factors were added 2 days after plating. The symbols represent the mean number of cells per field obtained from 4 cultures. Asterisks indicate differences that are significantly different from control ($P \leq 0.01$).

population stabilized. From day 7–27 *in vitro*, the number of neurons decreased by 81% (Fig. 1). Treatment of cultures with CNTF (10 ng/ml) resulted in a marked increase in the number of surviving neurons relative to control. This increase was statistically significant at 9, 13 and 16 days after plating ($P \leq 0.01$). Treatment of cells with BDNF (1 ng/ml) or bFGF (2 ng/ml) failed to produce significant changes in the survival of cells above those observed under control conditions. However, even in CNTF, viable neurons decreased during culture, and there was no significant difference in the number of neurons relative to control at times ≥ 20 days. Cells surviving through the first month were viable for the next three months in culture, as noted by previous investigators (Ransom et al., 1977; Keller et al., 1999). This pattern of cell survival was essentially the same in either MEM or Neurobasal media.




Fig. 2. Evoked release of neurotransmitters. Spinal cord cultures at 9 days post-plating were washed in HBSS, incubated with the appropriate radiolabeled neurotransmitters or precursors and stimulated in 56 mM K^+ in the presence of 2 mM Ca^{2+} . Cells were grown in MEM supplemented with N3 and 5% horse serum without additional growth factors. Bars represent the mean \pm SEM of 6 experiments.




Fig. 3. Time-dependent increase in glycine release. Evoked release of [³H]-glycine from cells treated with CNTF, bFGF or BDNF was measured at 7, 14 or 21 days after plating. There was a pronounced increase in the amount of [³H]-glycine released over this time period, especially between 14 and 21 days, although no consistent effect of the growth factors was observed. The cells were cultured in Neurobasal media supplemented with B27 and 2% horse serum. Bars represent the mean \pm SEM of 5 experiments. Asterisks indicate differences that are significantly different from [³H]-glycine released at 7 days ($P \leq 0.01$).

Depolarization- and Ca^{2+} -dependent neurotransmitter release

Cultures grown in MEM with N3 supplement at 9 days *in vitro* are capable of taking up and releasing approximately 5% of total incorporated glycine as indicated in Fig. 2. Uptake of either [³H]-glutamate or [³H]-glutamine was also observed and led to evoked release of 2–5% of the total incorporated label. However, although [¹⁴C]-Ch was taken up by these cultures, evoked release of labeled Ch/ACh was barely detectable (Fig. 2).

As cultures matured over a 3-week period, there was an increase in the percentage of [³H]-glycine released by K^+ -induced




Fig. 4. Time-dependent Ch/ACh release. Evoked release of Ch/ACh from cells treated with CNTF, bFGF and BDNF was measured at 7, 14 and 21 days after plating. Although there was an increase in the amount of Ch/ACh released in cultures treated with CNTF at 14 days, this effect was transient and no significant Ch/ACh release could be elicited at times ≥ 21 days. Cells were grown in MEM supplemented with N3 and 5% horse serum, but similar results were also observed with cells cultured in Neurobasal medium with B27 and 2% horse serum. Bars represent the mean \pm SEM of 5 experiments. Asterisk indicates significant difference from value at 7 days in culture ($P \leq 0.01$).

depolarization in the presence of normal extracellular Ca^{2+} . As illustrated in Fig. 3, this increase occurred predominantly between 14 and 21 days in culture and was not affected significantly by the presence of CNTF, bFGF or BDNF. Paradoxically, enhanced ^3H -glycine release occurred during the time when the total number of neurons in the cultures underwent marked reductions. Decrements in release expected from the loss of neurons may have been compensated by a continued expansion of dendritic processes from surviving cells (Ransom et al., 1977; Williamson et al., 1992). In addition, it is possible that fewer glycinergic neurons were lost during the time of rapid cell death, resulting in a higher percentage of glycinergic cells in the total population at day 21.

CNTF transiently increases Ch/ACh release

In contrast to the results obtained with ^3H -glycine release, evoked release of labeled Ch/ACh was generally undetectable in the absence of added CNTF (Fig. 4). This was found in both MEM and Neurobasal culture media and at all times up to 5 weeks after isolation of cells. However, even in CNTF, release of Ch/ACh was consistently observed only at two weeks in vitro. At this time, there were significant differences in the number of surviving neurons between control and CNTF-treated cultures, although the maximum difference occurred at day 7 (Fig. 1). Combinations of CNTF with BDNF or bFGF did not improve Ch/ACh release compared with treatment with CNTF alone (data not shown).

To establish that the 10 ng/ml CNTF used in this study was optimal, cultures were grown in the presence of 0.01 to 100 ng/ml CNTF for 12 days, starting on culture day 2, and assayed for high K^+ -evoked Ch/ACh release. Fig. 5 shows the effect of CNTF concentrations on release of radiolabeled Ch/ACh at the 2-week time window of optimum cholinergic response. There

was a significant increase in Ch/ACh release in the presence of CNTF concentrations >0.1 ng/ml, with maximal enhancement of release occurring at 10 ng/ml.

Discussion

There was a pronounced time-dependent decline in the number of neurons surviving in primary mouse spinal cord cultures, especially during the first two weeks after plating (Fig. 1), in agreement with previous investigators (Ransom et al., 1977; Bennett et al., 1980; Comella et al., 1994). Although the number of viable neurons in the cultures decreased, evoked release of neurotransmitters, notably glycine, increased during this time as indicated in Fig. 3. The increase in transmitter release parallels enhancements in neurite density, metabolic activity and electrical excitability (Ransom et al., 1977; Williamson et al., 1992). It is likely that the ability of the cultures to release neurotransmitters is dependent upon the development of an extensive field of neurites and synaptic connections. For release of glycine or glutamate, these factors more than compensate for the loss of neurons between initial isolation and development of a more mature culture. However, the selective loss of cholinergic neurons makes it difficult to elicit Ch/ACh release from spinal cord cultures in the absence of stabilization by coculture with target cells (Bennett et al., 1980; Kobayashi et al., 1987; Wang et al., 1990; Braun et al., 1997).

The presence of CNTF transiently delayed the death of viable neurons, with the maximum enhancement of cell survival occurring 9–16 days in vitro (Fig. 1). This enhanced survival was not detected in the presence of other neurotrophic factors such as BDNF or bFGF. Evoked release of Ch/ACh could be observed in the spinal cord cultures within a narrow window at 2 weeks in vitro (Fig. 4), in the presence of CNTF concentrations >0.1 ng/ml (Fig. 5). A plausible interpretation is that this restricted window represents the time between expansion of neurites and synaptic contacts, which increases overall evoked transmitter release, and the progressive deletion of cholinergic neurons, which results in the loss of Ch/ACh release. Although CNTF helped to stabilize the population of neurons up to day 16 in vitro, it could not prevent cell death at later time points. The reduction of Ch/ACh release in CNTF-treated cultures at ≥ 3 weeks is consistent with previous reports that cholinergic neurons represent a major population lost in cultures of embryonic spinal cord cells (Comella et al., 1994).

The finding that CNTF was effective in preserving evoked release of Ch/ACh, even transiently, was unexpected. Although CNTF has been implicated in maintaining motoneuron development in vivo (Forger et al., 1993; Masu et al., 1993), its efficacy in cultured cells has not previously been demonstrated (Hughes et al., 1993; Wong et al., 1993; Zurn et al., 1996). Of the growth factors examined, BDNF has been associated most often with enhanced motoneuron survival in vitro (Henderson et al., 1993; Hughes et al., 1993; Wong et al., 1993; Zurn et al., 1996; Kwon and Gurney, 1996). It should be noted that most of these in vitro studies examined periods only up to 5 days in culture. This would be too brief to show appreciable evoked neurotransmitter release in the assay employed in the current study.




Fig. 5. Effect of CNTF on Ch/ACh (♦) release. Cells were cultured in Neurobasal media supplemented with B27 and 2% horse serum. Cultures were treated with CNTF at the indicated concentrations starting on day 2. At 14 days in vitro, cells were loaded with ^{14}C -Ch, and evoked release in the presence of 54 mM K^+ and 2 mM Ca^{2+} was determined as a percent of total labeled transmitter. CNTF increased ^{14}C -Ch/ACh release in a concentration-dependent manner with 10 ng/ml being optimal. Symbols represent the mean \pm SEM of data from 6 experiments. Asterisks indicate differences that are significantly different from control values ($P \leq 0.01$).

Although CNTF was found to be beneficial in preserving cholinergic function, the brief window of cholinergic neurotransmission would limit the nature of studies that could be undertaken. A more stable response would be desirable to study problems such as BoNT persistence in cholinergic motoneuron terminals that represent their primary *in vivo* target. Perhaps a combination of treatment by CNTF and coculture with potential targets such as skeletal muscle may lead to the required stability of cholinergic responses.

References

- Bennett, M.R., Lai, K., Nurcombe, V., 1980. Identification of embryonic motoneurons *in vitro*: their survival is dependent on skeletal muscle. *Brain Research* 190 (2), 537–542.
- Bergey, G.K., Bigalke, H., Nelson, P.G., 1987. Differential effects of tetanus toxin on inhibitory and excitatory synaptic transmission in mammalian spinal cord neurons in culture: a presynaptic locus of action for tetanus toxin. *Journal of Neurophysiology* 57 (1), 121–131.
- Braun, S., Croizat, B., Lagrange, M.C., Poindron, P., Warter, J.M., 1997. Degeneration of cocultures of spinal muscular atrophy muscle cells and rat spinal cord explants is not due to secreted factors and cannot be prevented by neurotrophins. *Muscle and Nerve* 20 (8), 953–960.
- Brenneman, D.E., Neale, E.A., Foster, G.A., d'Autremont, S.W., Westbrook, G.L., 1987. Nonneuronal cells mediate neurotrophic action of vasoactive intestinal peptide. *The Journal of Cell Biology* 104 (6), 1603–1610.
- Brewer, G.J., 1995. Serum-free B27/neurobasal medium supports differentiated growth of neurons from the striatum, substantia nigra, septum, cerebral cortex, cerebellum, and dentate gyrus. *Journal of Neuroscience Research* 42 (5), 674–683.
- Comella, J.X., Sanz-Rodriguez, C., Aldea, M., Esquerda, J.E., 1994. Skeletal muscle-derived trophic factors prevent motoneurons from entering an active cell death program *in vitro*. *The Journal of Neuroscience* 14 (5), 2674–2686.
- Daniels-Holgate, P.U., Dolly, J.O., 1996. Productive and non-productive binding of botulinum neurotoxin A to motor nerve endings are distinguished by its heavy chain. *Journal of Neuroscience Research* 44 (3), 263–271.
- Forger, N.G., Roberts, S.L., Wong, V., Breedlove, S.M., 1993. Ciliary neurotrophic factor maintains motoneurons and their target muscles in developing rats. *The Journal of Neuroscience* 13 (11), 4720–4726.
- Hall, Y.H., Chaddock, J.A., Moulds, H.J., Kirby, E.R., Alexander, F.C., Marks, J.D., Foster, K.A., 2004. Novel application of an *in vitro* technique to the detection and quantification of botulinum neurotoxin antibodies. *Journal of Immunological Methods* 288 (1–2), 55–60.
- Henderson, C.E., Camu, W., Mettling, C., Gouin, A., Poulsen, K., Karihaloo, M., Rullamas, J., Evans, T., McMahon, S.B., Armanini, M.P., Berkemeier, L., Phillips, H.S., Rosenthal, A., 1993. Neurotrophins promote motor neuron survival and are present in embryonic limb bud. *Nature* 363 (6426), 266–270.
- Hughes, R.A., Sendtner, M., Thoenen, H., 1993. Members of several gene families influence survival of rat motoneurons *in vitro* and *in vivo*. *Journal of Neuroscience Research* 36 (6), 663–671.
- Keller, J.E., Neale, E.A., Oyler, G., Adler, M., 1999. Persistence of botulinum neurotoxin actions in cultured spinal cord cells. *FEBS Letters* 456 (1), 137–142.
- Keller, J.E., Cai, F., Neale, E.A., 2004. Uptake of botulinum neurotoxin into cultured neurons. *Biochemistry* 43 (2), 526–532.
- Kobayashi, T., Askanas, V., Engel, W.K., 1987. Human muscle cultured in monolayer and cocultured with fetal rat spinal cord: importance of dorsal root ganglia for achieving successful functional innervation. *The Journal of Neuroscience* 7 (10), 3131–3141.
- Kwon, Y.W., Gurney, M.E., 1996. Brain-derived neurotrophic factor transiently stabilizes silent synapses on developing neuromuscular junctions. *Journal of Neurobiology* 29 (4), 503–516.
- Masu, Y., Wolf, E., Holtmann, B., Sendtner, M., Brem, G., Thoenen, H., 1993. Disruption of the CNTF gene results in motor neuron degeneration. *Nature* 365 (6441), 27–32.
- McGee, R., Simpson, P., Christian, C., Mata, M., Nelson, P., Nirenberg, M., 1978. Regulation of acetylcholine release from neuroblastoma x glioma hybrid cells. *Proceedings of the National Academy of Sciences* 75 (3), 1314–1318.
- Ransom, B.R., Neale, E., Henkart, M., Bullock, P.N., Nelson, P.G., 1977. Mouse spinal cord in cell culture. I. Morphology and intrinsic neuronal electrophysiologic properties. *Journal of Neurophysiology* 40 (5), 1132–1150.
- Romijn, H.J., Habets, A.M., Mud, M.T., Wolters, P.S., 1981. Nerve outgrowth, synaptogenesis and bioelectric activity in fetal rat cerebral cortex tissue cultured in serum-free, chemically defined medium. *Brain Research* 254 (4), 583–589.
- Sheridan, R.E., Smith, T.J., Adler, M., 2005. Primary cell culture for evaluation of botulinum neurotoxin antagonists. *Toxicon* 45 (3), 377–382.
- Simpson, L.L., 2004. Identification of the major steps in botulinum toxin action. *Annual Review of Pharmacology and Toxicology* 44, 167–193.
- Wang, F.Z., Nelson, P.G., Fitzgerald, S.C., Hersh, L.B., Neale, E.A., 1990. Cholinergic function in cultures of mouse spinal cord neurons. *Journal of Neuroscience Research* 25 (3), 312–323.
- Williamson, L.C., Fitzgerald, S.C., Neale, E.A., 1992. Differential effects of tetanus toxin on inhibitory and excitatory neurotransmitter release from mammalian spinal cord cells in culture. *Journal of Neurochemistry* 59 (6), 2148–2157.
- Williamson, L.C., Halpern, J.L., Montecucco, C., Brown, J.E., Neale, E.A., 1996. Clostridial neurotoxins and substrate proteolysis in intact neurons: botulinum neurotoxin C acts on synaptosomal-associated protein of 25 kDa. *Journal of Biological Chemistry* 271 (13), 7694–7699.
- Wong, V., Arriaga, R., Ip, N.Y., Lindsay, R.M., 1993. The neurotrophins BDNF, NT-3 and NT-4/5, but not NGF, up-regulate the cholinergic phenotype of developing motor neurons. *European Journal of Neuroscience* 5 (5), 466–474.
- Zurn, A.D., Winkel, L., Menoud, A., Djabali, K., Aebischer, P., 1996. Combined effects of GDNF, BDNF, and CNTF on motoneuron differentiation *in vitro*. *Journal of Neuroscience Research* 44 (2), 133–141.



R00118203_LFS_11241